

The Isomorphism Conjectures in the torsion free case (Lecture II)

Wolfgang Lück

Bonn

Germany

email wolfgang.lueck@him.uni-bonn.de

<http://131.220.77.52/lueck/>

Berlin, June 19, 2012

Flashback

- We introduced $K_n(R)$ for $n \in \mathbb{Z}$.
- We discussed the topological relevance of $K_0(RG)$ and the Whitehead group $\text{Wh}(G)$, e.g., **the finiteness obstruction** and the **s -cobordism theorem**.
- We stated the conjectures that $\tilde{K}_0(\mathbb{Z}G)$ and $\text{Wh}(G)$ vanish for torsion free G .
- We presented the **Bass-Heller-Swan decomposition** and indicated some similarities between $K_n(RG)$ and **group homology**.
- **Cliffhanger**

Question (K -theory of group rings and group homology)

Is there a relationship between $K_n(RG)$ and the group homology of G ?

- We introduce **spectra** and how they yield **homology theories**.
- We state the **Farrell-Jones Conjecture** and the **Baum-Connes Conjecture** for torsion free groups.
- We discuss applications of these conjectures, such as the **Kaplansky Conjecture** and the **Borel Conjecture**.
- We explain that the formulations for torsion free groups cannot extend to arbitrary groups.

Definition (Spectrum)

A *spectrum*

$$\mathbf{E} = \{(E(n), \sigma(n)) \mid n \in \mathbb{Z}\}$$

is a sequence of pointed spaces $\{E(n) \mid n \in \mathbb{Z}\}$ together with pointed maps called *structure maps*

$$\sigma(n): E(n) \wedge S^1 \longrightarrow E(n+1).$$

- Given two pointed spaces $X = (X, x_0)$ and $Y = (Y, y_0)$, their **one-point-union** and their **smash product** are defined to be the pointed spaces

$$X \vee Y := \{(x, y_0) \mid x \in X\} \cup \{(x_0, y) \mid y \in Y\} \subseteq X \times Y;$$

$$X \wedge Y := (X \times Y)/(X \vee Y).$$

- If X is a pointed space and \mathbf{E} is a spectrum, then we obtain a new spectrum by $X \wedge \mathbf{E}$.

Exercise

Show $S^{n+1} \cong S^n \wedge S^1$.

Definition (Homotopy groups of a spectrum)

Given a spectrum \mathbf{E} , define for $n \in \mathbb{Z}$ its *n -th homotopy group*

$$\pi_n(\mathbf{E}) := \operatorname{colim}_{k \rightarrow \infty} \pi_{k+n}(E(k))$$

to be the abelian group which is given by the colimit over the directed system indexed by \mathbb{Z}

$$\dots \xrightarrow{\sigma(k-1)_*} \pi_{k+n}(E(k)) \xrightarrow{\sigma(k)_*} \pi_{k+n+1}(E(k+1)) \xrightarrow{\sigma(k+1)_*} \dots$$

- Notice that a spectrum, in contrast to a space, can have non-trivial negative homotopy groups.

- Algebraic K -theory spectrum

For a ring R , there is the algebraic K -theory spectrum \mathbf{K}_R with the property

$$\pi_n(\mathbf{K}_R) = K_n(R) \quad \text{for } n \in \mathbb{Z}.$$

- Algebraic L -theory spectrum

For a ring with involution R , there is the algebraic L -theory spectrum $\mathbf{L}_R^{\langle -\infty \rangle}$ with the property

$$\pi_n(\mathbf{L}_R^{\langle -\infty \rangle}) = L_n^{\langle -\infty \rangle}(R) \quad \text{for } n \in \mathbb{Z}.$$

Definition (Homology theory)

A *homology theory* \mathcal{H}_* is a covariant functor from the category of CW-pairs to the category of \mathbb{Z} -graded abelian groups together with natural transformations

$$\partial_n(X, A): \mathcal{H}_n(X, A) \rightarrow \mathcal{H}_{n-1}(A)$$

for $n \in \mathbb{Z}$ satisfying the following axioms:

- Homotopy invariance;
- Long exact sequence of a pair;
- Excision;
- Disjoint union axiom.

Theorem (Homology theories and spectra)

Let \mathbf{E} be a spectrum.

Then we obtain a homology theory $H_*(-; \mathbf{E})$ by

$$H_n(X, A; \mathbf{E}) := \pi_n((X \cup_A \text{cone}(A)) \wedge \mathbf{E}).$$

It satisfies

$$H_n(\text{pt}; \mathbf{E}) = \pi_n(\mathbf{E}).$$

The Isomorphism Conjectures for torsion free groups

Conjecture (*K*-theoretic Farrell-Jones Conjecture for torsion free groups and regular rings)

The *K*-theoretic Farrell-Jones Conjecture with coefficients in the regular ring R for the torsion free group G predicts that the *assembly map*

$$H_n(BG; \mathbf{K}_R) \rightarrow K_n(RG)$$

is bijective for every $n \in \mathbb{Z}$.

- $K_n(RG)$ is the algebraic K -theory of the group ring RG ;
- \mathbf{K}_R is the (non-connective) algebraic K -theory spectrum of R ;
- $H_n(\text{pt}; \mathbf{K}_R) \cong \pi_n(\mathbf{K}_R) \cong K_n(R)$ for $n \in \mathbb{Z}$.
- BG is the *classifying space* of the group G .

Conjecture (*L*-theoretic Farrell-Jones Conjecture for torsion free groups)

The *L*-theoretic Farrell-Jones Conjecture with coefficients in the ring with involution R for the torsion free group G predicts that the assembly map

$$H_n(BG; \mathbf{L}_R^{\langle -\infty \rangle}) \rightarrow L_n^{\langle -\infty \rangle}(RG)$$

is bijective for every $n \in \mathbb{Z}$.

- $L_n^{\langle -\infty \rangle}(RG)$ is the algebraic L -theory of RG with decoration $\langle -\infty \rangle$;
- $\mathbf{L}_R^{\langle -\infty \rangle}$ is the algebraic L -theory spectrum of R with decoration $\langle -\infty \rangle$;
- $H_n(\text{pt}; \mathbf{L}_R^{\langle -\infty \rangle}) \cong \pi_n(\mathbf{L}_R^{\langle -\infty \rangle}) \cong L_n^{\langle -\infty \rangle}(R)$ for $n \in \mathbb{Z}$.

Conjecture (Baum-Connes Conjecture for torsion free groups)

The *Baum-Connes Conjecture* for the torsion free group predicts that the *assembly map*

$$K_n(BG) \rightarrow K_n(C_r^*(G))$$

is bijective for every $n \in \mathbb{Z}$.

- $K_n(BG)$ is the topological K -homology of BG .
- $K_n(C_r^*(G))$ is the topological K -theory of the reduced complex group C^* -algebra $C_r^*(G)$ of G .

Exercise

Let G be the fundamental group of a closed orientable 2-manifold. Compute $K_n(BG)$.

Conclusions of the Isomorphism Conjectures for torsion free groups

- In order to illustrate the depth of the Farrell-Jones Conjecture and the Baum-Connes Conjecture, we present some conclusions which are interesting in their own right.
- Let $\mathcal{FJ}_K(R)$, respectively $\mathcal{FJ}_L(R)$, be the class of groups that satisfy the K -theoretic, respectively L -theoretic, Farrell-Jones Conjecture for the coefficient ring R .
- Let \mathcal{BC} be the class of groups that satisfy the Baum-Connes Conjecture.

Lemma

Let R be a regular ring. Suppose that G is torsion free and $G \in \mathcal{FJ}_K(R)$. Then

- $K_n(RG) = 0$ for $n \leq -1$;
- The change of rings map $K_0(R) \rightarrow K_0(RG)$ is bijective. In particular $\tilde{K}_0(RG)$ is trivial if and only if $\tilde{K}_0(R)$ is trivial.

Lemma

Suppose that G is torsion free and $G \in \mathcal{FJ}_K(\mathbb{Z})$. Then the Whitehead group $\text{Wh}(G)$ is trivial.

Proof.

The idea of the proof is to study the **Atiyah-Hirzebruch spectral sequence** converging to $H_n(BG; \mathbf{K}_R)$ whose E^2 -term is given by

$$E_{p,q}^2 = H_p(BG, K_q(R)).$$



In particular, for a torsion free group $G \in \mathcal{FJ}_K(\mathbb{Z})$ we get:

- $K_n(\mathbb{Z}G) = 0$ for $n \leq -1$;
- $\tilde{K}_0(\mathbb{Z}G) = 0$;
- $\text{Wh}(G) = 0$;

- Every finitely dominated CW -complex X with $G = \pi_1(X)$ is homotopy equivalent to a finite CW -complex;

- Every compact h -cobordism W of dimension ≥ 6 with $\pi_1(W) \cong G$ is trivial;

- If G belongs to $\mathcal{FJ}_K(\mathbb{Z})$, then it is of type FF if and only if it is of type FP (**Serre's problem**).

Mathematicians!



What my parents think I do



What my friends think I do



What my students think I do




What my spouse thinks I do



What my colleagues think I do



What I actually do





Conjecture (Kaplansky Conjecture)

The *Kaplansky Conjecture* says that for a torsion free group G and an integral domain R the elements 0 and 1 are the only idempotents in RG .

Theorem (The Farrell-Jones Conjecture and the Kaplansky Conjecture)

Let F be a skew-field and let G be a group with $G \in \mathcal{FJ}_K(F)$. Suppose that one of the following conditions is satisfied:

- F is commutative and has characteristic zero, and G is torsion free;
- G is torsion free and sofic;
- The characteristic of F is p , all finite subgroups of G are p -groups and G is sofic;

Then 0 and 1 are the only idempotents in FG .

Proof.

- We only treat the case of fields of characteristic zero.
- Let p be an idempotent in FG . We want to show $p \in \{0, 1\}$.
- Denote by $\epsilon: FG \rightarrow F$ the augmentation homomorphism sending $\sum_{g \in G} r_g \cdot g$ to $\sum_{g \in G} r_g$. It suffices to show $p = 0$ under the assumption that $\epsilon(p) = 0$.
- Let $(p) \subseteq FG$ be the ideal generated by p , which is a finitely generated projective FG -module.
- Since $G \in \mathcal{FJ}_K(F)$, we can conclude that

$$i_*: K_0(F) \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow K_0(FG) \otimes_{\mathbb{Z}} \mathbb{Q}$$

is surjective.

- Hence we can find a finitely generated projective F -module P and integers $k, m, n \geq 0$ satisfying

$$(p)^k \oplus FG^m \cong_{FG} i_*(P) \oplus FG^n.$$

Proof (continued).

- If we now apply $i_* \circ \epsilon_*$ and use $\epsilon \circ i = \text{id}$, $i_* \circ \epsilon_*(FG^l) \cong FG^l$ and $\epsilon(p) = 0$, then we obtain

$$FG^m \cong i_*(P) \oplus FG^n.$$

- Inserting this in the first equation yields

$$(p)^k \oplus i_*(P) \oplus FG^n \cong i_*(P) \oplus FG^n.$$

- Our assumptions on F and G imply that FG is **stably finite**, i.e., if A and B are square matrices over FG with $AB = I$, then $BA = I$. This implies $(p)^k = 0$ and hence $p = 0$.



Conjecture (Borel Conjecture)

*The **Borel Conjecture for G** predicts that for two closed aspherical manifolds M and N with $\pi_1(M) \cong \pi_1(N) \cong G$ any homotopy equivalence $M \rightarrow N$ is homotopic to a homeomorphism and in particular that M and N are homeomorphic.*

- In particular the Borel Conjecture predicts that two closed aspherical manifolds are homeomorphic if and only if their fundamental groups are isomorphic.

- The Borel Conjecture can be viewed as the topological version of **Mostow rigidity**.

A special case of Mostow rigidity says that any homotopy equivalence between closed hyperbolic manifolds of dimension ≥ 3 is homotopic to an isometric diffeomorphism.

- The Borel Conjecture is not true in the smooth category by results of **Farrell-Jones**.
- There are also non-aspherical manifolds that are topologically rigid in the sense of the Borel Conjecture (see **Kreck-L.**).

Theorem (The Farrell-Jones Conjecture and the Borel Conjecture)

If the K - and L -theoretic Farrell-Jones Conjecture hold for G in the case $R = \mathbb{Z}$, then the Borel Conjecture is true in dimension ≥ 5 and in dimension 4 if G is good in the sense of Freedman.

- **Thurston's Geometrization Conjecture** implies the Borel Conjecture in dimension 3.

Exercise

Prove the Borel Conjecture in dimensions 1 and 2.

Definition (Structure set)

The *structure set* $S^{\text{top}}(M)$ of a manifold M consists of equivalence classes of orientation preserving homotopy equivalences $N \rightarrow M$ with a manifold N as source.

Two such homotopy equivalences $f_0: N_0 \rightarrow M$ and $f_1: N_1 \rightarrow M$ are equivalent if there exists a homeomorphism $g: N_0 \rightarrow N_1$ with $f_1 \circ g \simeq f_0$.

Theorem

The Borel Conjecture holds for a closed manifold M if and only if $S^{\text{top}}(M)$ consists of one element.

Theorem (Ranicki)

There is an exact sequence of abelian groups, called *the algebraic surgery exact sequence*, for an n -dimensional closed manifold M

$$\begin{array}{ccccccc} \dots & \xrightarrow{\sigma_{n+1}} & H_{n+1}(M; \mathbf{L}\langle 1 \rangle) & \xrightarrow{A_{n+1}} & L_{n+1}(\mathbb{Z}\pi_1(M)) & \xrightarrow{\partial_{n+1}} & \\ & & & & \mathcal{S}^{\text{top}}(M) & \xrightarrow{\sigma_n} & H_n(M; \mathbf{L}\langle 1 \rangle) & \xrightarrow{A_n} & L_n(\mathbb{Z}\pi_1(M)) & \xrightarrow{\partial_n} & \dots \end{array}$$

It can be identified with the classical geometric surgery sequence due to *Sullivan and Wall* in high dimensions.

- $\mathcal{S}^{\text{top}}(M)$ consists of one element if and only if A_{n+1} is surjective and A_n is injective.
- $H_k(M; \mathbf{L}\langle 1 \rangle) \rightarrow H_k(M; \mathbf{L})$ is bijective for $k \geq n + 1$ and injective for $k = n$.

What happens for groups with torsion?

- The versions of the Farrell-Jones Conjecture and the Baum-Connes Conjecture above are false for finite groups unless the group is trivial.
- For instance the version of the Baum-Connes Conjecture above would predict that for a finite group G

$$K_0(BG) \cong K_0(C_r^*(G)) \cong R_{\mathbb{C}}(G).$$

However, $K_0(BG) \otimes_{\mathbb{Z}} \mathbb{Q} \cong_{\mathbb{Q}} K_0(\text{pt}) \otimes_{\mathbb{Z}} \mathbb{Q} \cong_{\mathbb{Q}} \mathbb{Q}$ and $R_{\mathbb{C}}(G) \otimes_{\mathbb{Z}} \mathbb{Q} \cong_{\mathbb{Q}} \mathbb{Q}$ holds if and only if G is trivial.

- If G is torsion free, then the version of the K -theoretic Farrell-Jones Conjecture predicts

$$\begin{aligned} H_n(B\mathbb{Z}; \mathbf{K}_R) &= H_n(S^1; \mathbf{K}_R) = H_n(\text{pt}; \mathbf{K}_R) \oplus H_{n-1}(\text{pt}; \mathbf{K}_R) \\ &= K_n(R) \oplus K_{n-1}(R) \cong K_n(R\mathbb{Z}). \end{aligned}$$

In view of the Bass-Heller-Swan decomposition this is only possible if $NK_n(R)$ vanishes which is true for regular rings R but not for general rings R .

Question (Arbitrary groups and rings)

Are there versions of the Farrell-Jones Conjecture for arbitrary groups and rings and of the Baum-Connes Conjecture for arbitrary groups?

To be continued

Stay tuned

Next talk: Tomorrow at 14:00